

CIR-191776/2021-Imidacloprid (429)-3930

Imidacloprid 2.15% Gel

(Household Insecticide)

It contains Imidacloprid 2.15% as gel.

Recommendation

This gel is recommended for effective control of German and American cockroaches in houses. Cockroaches are attracted to the gel and after feeding, the cockroaches are killed. Application of gel is convenient for use.

Direction of Use

Remove the cap from the cartridge tip and fit the needle or use directly to apply gel. Touch the tip to the surface and depress the plunger of the cartridge or lever of the bait gun. Gel is applied as spots or as a line along cracks and crevices or places where cockroaches are like to hide or breed. For an effective control, apply 3 to 4 gel spots/sq.m area and each gel spot with 3-5 m diameter in size (approximately 0.1g/spot). If cockroaches are seen, reapply gel. Do not apply gel in areas which are frequently washed.

Precaution

Keep out of reach of children and pests. Avoid contact with skin and clothing. Keep exposed gel away from open foods and food contact surfaces. Wash hands after handling. Do not store or expose gel in high temperature areas. do not smoke or chew pan or tobacco while handling the product. Wash exposed body parts with soap and water after handling and before eating, drinking and smoking.

Symptoms Of Poisoning

Apathy, Myatonia, Tremor, difficult breathing & myospasms.

First Aid

Induce vomiting if swallowed by thickling at the back of the throat with the finger. For eye contact, flush eyes with plenty of cold water. If contaminated with skin, wash skin with plenty of soap and water.

<u>Antidote</u>

No specific antidote is known. Treat symptomatically.

Disposal Of Used Container

Destroy empty cartridges by crushing and burying in soil in a safe place away from water sources and habitants. Used cartridges should be destroyed in such a manner so that there will be no water or environmental pollution. Used cartridges should not be left outside to prevent its reuse.



Storage Conditions

The cartridge containing the gel should in original containers in a separate room or almirah under lock and key depending on the quantity of insecticide, away from reach of children, food stuffs, animal feeds and other articles and keep in cool and dry place. The premises for storage should be well built, well lit, sufficient in dimension and well ventilated.

Chemical Composition:

	Total: 100.000 % w/w
Distilled water	Q.S %
disulfonate)	
(dioxoin dan-2yl) quinoline	
Colourant : Yellow (Sodium 2-	0.10 % w/w
Gelling agent : Seaweed extract	2.50 % w/w
Stabiliser : BHT	0.25 % w/w
Thickener: Polysaccharide gum	0.20 % w/w
Humectant : Glycerine	10.00 % w/w
Phagostimulant : Fruit Sugar (Levulose)	55.00 % w/w
Active Ingredient : Imidacloprid	2.15 % w/w